



(19)

Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 890 639 A2

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:

13.01.1999 Bulletin 1999/02

(51) Int. Cl.⁶: **C12N 15/12, C07K 14/47**

(21) Application number: **98112742.6**

(22) Date of filing: **09.07.1998**

(84) Designated Contracting States:

**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE**

Designated Extension States:

AL LT LV MK RO SI

(30) Priority: **09.07.1997 EP 97111602**

(71) Applicant:

**Gesellschaft für biotechnologische
Forschung mbH (GBF)
38124 Braunschweig (DE)**

(72) Inventors:

- **Ahrens, Marion**
38124 Braunschweig (DE)
- **Bächner, Dietmar**
38124 Braunschweig (DE)

• **Hoffmann, Andrea**

38124 Braunschweig (DE)

• **Lauber, Jörg**

38124 Braunschweig (DE)

• **Steinert, Peter**

38124 Braunschweig (DE)

• **Flohé, Leopold**

38124 Braunschweig (DE)

• **Gross, Claus-Gerhard**

38124 Braunschweig (DE)

(74) Representative:

Boeters, Hans Dietrich, Dr. et al

• **Patentanwälte Boeters & Bauer,**

Bereiteranger 15

81541 München (DE)

(54) **BMP2-induced cDNA and its use**

(57) The recombinant expression of human BMP2, BMP4 - BMP7 in murine mesenchymal C3H10T $\frac{1}{2}$ progenitors mediates differentiation into three mesenchymal lineages in different efficiencies: the osteogenic, the chondrogenic and the adipogenic lineage. This developmental *in vitro* model was used to identify and to characterize cDNAs involved during the manifestation of these lineages *in vivo*. By subtractive cloning an as yet undescribed cDNA, 29A has been cloned which encodes a putative secreted factor which is expressed in developing osteo- /chondrogenic tissues of vertebrae, ribs, tooth and the limb bud.

EP 0 890 639 A2

Description

Introduction

During embryonic development, the vertebrate skeletal elements arise from mesenchymal cell-condensations forming cartilage or bone, eventually. Recent studies suggest that bone morphogenetic proteins (BMPs) may play a crucial role during the formation of skeletal condensations and the onset of differentiation from mesenchymal progenitors. The mammalian bone morphogenetic proteins (BMPs) were originally purified and characterized from adult bone on the basis of their ability to induce a cascade of events leading to ectopic bone formation if implanted subcutaneously or at intra-muscular sites.

Molecular cloning of these genes and biochemical characterizations established BMPs (with the exception of BMP1) as a family of proteins which are generated from dimeric precursors proteolytically processed to 25,000-30,000 M_r homo- or heterodimers belonging to the transforming growth factor β (TGF- β) superfamily. Members of this family can be classified to the degree of amino acid identity of their C-terminal domains. The BMPs also share a high identity to other closely related proteins which have been characterized in *Xenopus* and *Drosophila* like the *decapentaplegic* (*dpp*) gene, the latter being involved both in dorsoventral body patterning and in imaginal disk formation (Irish and Gelbart, 1987; Ferguson and Anderson, 1992). The BMPs are also related to Vgl, which in *Xenopus* has been postulated to play a role in embryonic development and mesoderm specification (Blessing et al., 1993; Weeks and Melton, 1987; Lyons et al., 1989b; Lyons et al., 1989a). While the homogenous inactivation of the murine BMP4 as well as the BMP type Ia receptor genes results in an early embryonic lethality consistent with a putative role in early mesoderm formation (Winner et al., 1995; Mishina et al., 1995), BMP7 null mice exhibit only relatively mild skeletal abnormalities affecting mesenchymal condensations rather than chondrogenic differentiation (Karsenty et al., 1996; Hofmann et al., 1996).

BMPs induce condensations and chondrogenesis in primary cells and cell lines derived from limb buds. The potency of the various BMPs (BMP2-7) differ in these primary systems but in all cases they directly mediate chondrocytic differentiation.

The murine fibroblastic C3H10T $\frac{1}{2}$ cell line which has been established from an early-stage mouse embryo represents a relatively early stage of mesenchymal cell determination with the ability to differentiate into myoblasts, adipocytes, chondrocytes and osteoblasts (Reznikoff et al., 1973; Taylor and Jones, 1979; Wang et al., 1993; Ahrens et al., 1993). Their responsiveness towards TGF- β and BMP-treatment make this line a useful model system to explore the involvement of factors in various mesenchymal differentiation processes. BMP2 and BMP4 possess the ability in mesenchymal progenitor C3H10T $\frac{1}{2}$ cells to mediate the differentiation into chondrocytes, osteoblasts and adipocytes (Wang et al., 1993; Ahrens et al., 1993). The extension of the analysis onto other members of the family BMP5-7 in this study shows that all BMPs investigated possess the potency to mediate differentiation into three mesenchymal cell types, albeit in largely varying efficiencies. To assess the value of the C3H10T $\frac{1}{2}$ *in vitro* model as a model for BMP-mediated mesenchymal differentiation we performed subtractive cloning analysis for BMP-upregulated genes in this system and characterized their expression in the embryonic development in the mouse by *in situ* hybridization. We were able to isolate BMP-upregulated genes typical for the osteo- /chondrogenic and adipogenic development. In addition we characterized BMP-upregulated genes and cDNAs which are unknown or as yet have not been assigned to BMP-mediated mesenchymal development. Among these genes is a new putative secreted factor 29A which is expressed in osteoblasts, in murine embryonic development in precartilaginous condensation and during limb and tooth development.

Materials and Methods

Cell Lines, culture conditions and transfection experiments

The features of BMP2 and BMP4 transfected C3H10T $\frac{1}{2}$ cells have been described by Ahrens et al. (1993). Human BMP2 and BMP4 are constitutively transcribed by the LTR of the myeloproliferative sarcoma virus (MPSV). Human versions of BMP5, BMP6 and BMP7 have been described in Wozney et al. (1989) and incorporated in the expression vector described before. Transfection was performed by calcium-phosphate precipitation. Control or BMP-transfected C3H10T $\frac{1}{2}$ cells were selected by cotransfection with pSV2pac mediating resistance against puromycin (5 μ g/ml). Puromycin-resistant colonies were subcultivated and selection-pressure was maintained during the entire cultivation period to follow. If not stated otherwise cells were plated at a density of 5,000 cells /cm². Cells were routinely grown in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% fetal calf serum. After reaching confluence (arbitrarily termed day 0) 50 μ g/ml ascorbic acid and 10 mM β -glycerolphosphate were added as specified in the protocol of Owen et al., (1990) for the cultivation of native osteoblast-like cells. Primary osteoblast-like cells were isolated from the calvariae of 5 day old mice (NMRI) by sequential collagenase digestion and cultivated as described (Owen et al., 1990).

mRNA analysis

C3H10T $\frac{1}{2}$ cells harboring the expression vectors were cultivated as described above. Cells were harvested at the indicated time intervals and total RNA was isolated by guanidinium/CsCl step-gradients. Total cellular RNA (10 μ g) was separated electrophoretically in a 2.2 M formaldehyde - 1.2% agarose gel and transferred to nitrocellulose. Hybridization was carried out with nick-translated 32 P-labelled gene-specific DNA probes.

Histological methods and verification of cellular phenotypes

Osteoblasts exhibit a stellate morphology displaying high levels of alkaline phosphatase activity which was visualized by cellular staining with α -naphthyl-phosphate and Fast Red. Deposition of mineral was monitored by von Kossa-staining method (Owen et al., 1990). Osteogenesis was also investigated by northern analyses with hybridization probes typical or specific for the osteogenic developmental sequence namely collagen I, osteopontin, osteonectin and osteocalcin. In addition, the up-regulation of the parathyroid hormone receptor (PTH/PTHrP-receptor) at the onset of the osteogenic developmental sequence was studied by northern analyses. Adipocytes were identified morphologically as oil-droplet filled cells and/or by staining with Oil Red O. Marker genes for adipocytes were the lipoprotein lipase and ap2. Chondrocytes were identified by staining with Alcian Blue at pH 2.5. They also displayed alkaline phosphatase activity but in comparison with osteoblasts differed in intensity and a rounded cell morphology. Marker genes for chondrocytes was collagen II.

5'- and 3'- extension of cDNA clones by the RACE - technology (Rapid Amplification of cDNA nds)

To complete the 29A cDNA 5'-RACE was performed essentially as described (Frohmann, 1990). Random primers were removed from the first strand cDNA preparation by gel purification (Jetsorb-Kit, Genomed), and a dA-tail was added to the 3'-end of the single-stranded DNA by terminal transferase in a 10 μ l-reaction containing 200 mM potassium cacodylate, 25 mM Tris pH 6.6, 0.25 mg/ml bovine serum albumin, 1.5 mM CoCl $_2$, 0.5 mM dATP and 10 U enzyme (25 min, 37°C). After phenolization and precipitation, the first PCR-amplification was performed using the primers R $_0$ -R $_1$ -(dT $_{17}$) and 3'-reverse primer (30 pmol each) under the same temperature conditions as detailed for RT-PCR except that 50°C was used for annealing. The region corresponding to the desired band was eluted from a low-melting agarose gel and reamplified using the primers R $_0$ and a new nested 3'- reverse primer (N $_1$; 30 pmol each). For the 3'-RACE, total RNA was reverse transcribed using 2.5 μ g Oligo-dT $_{12-18}$ (Gibco-BRL) as described above. Double-stranded cDNA was then immediately synthesized by adding 30 pmol 5'-primer and fresh dNTPs to the reaction mixture (1h, 37°C). The sample was gel-purified and the first PCR-amplification was performed using the primers R $_0$ -R $_1$ -(dT $_{17}$) and a new nested 5'-primer (N $_2$; 30 pmol each) followed by a second PCR with the primers R $_0$ and

Subtractive cloning

Subtractive cDNA cloning was performed after a modified protocol. *mRNA preparation.* BMP-2 and untransfected C3H10T $\frac{1}{2}$ cells were cultivated as described above. Eight days after reaching confluence cells were harvested and total RNA was isolated by guanidinium/CsCl step gradients. mRNA was purified from total RNA by oligo(dT)-cellulose chromatography. *cDNA synthesis and library construction.* mRNA (2.5 μ g) from BMP-2-transfected (target) and control C3H10T $\frac{1}{2}$ cells (driver) were reverse transcribed with Moloney murine leukemia virus reverse transcriptase (BRL) in the presence of 1 μ g of random hexanucleotide (Pharmacia). Second strand synthesis was performed with a commercially available cDNA Synthesis Kit (BRL). cDNA was degraded to ~500 bp by a short ultrasonic pulse (3 x 5 sec) in a Branson sonifier (250 watts). The latter step is thought to prevent a PCR-dependent amplification of short cDNA sequences in the subtractive cloning procedure. After sonification staggered ends were filled up by treatment with T4-DNA polymerase. The driver and the target cDNAs were ligated to different oligonucleotides harboring a *Eco* RI restriction or a *Hind* III restriction site, respectively (Duguid and Dinauer (1989)). For the driver cDNAs the sequence of the oligonucleotides was:

3' ATCAGGCTTAAGTTCGTTCTC 5'

5' TAGTCCGAATTCAGCAAGAGCACA 3'.

For the target library the sequence of oligonucleotides was:

3' TAGCAGTTCGAAGTTCATCG 5'

5' ATCGTCAAGCTTCAAGTTAGCATCG 3'.

cDNAs were amplified by PCR (30 cycles) using the different oligonucleotides as primers. The PCR-amplification of the control cDNA was performed with 0.25 mM Biotin-4-dUTP as dTTP analog. The PCR-products were purified with *Strataclean* Resin from *Stratagene* and the incorporation of the biotinylated nucleotides was verified by the *BluGENE* nonradioactive nucleic acid detection system from BRL. The amplified cDNA libraries from both control and BMP-2-

transfected C3H10T $\frac{1}{2}$ cells represent the starting material for the subtractive cloning procedure. *Library subtraction.* 15 μ g of the biotinylated control library cDNA was mixed with 1.5 μ g of the BMP-2-transfected library cDNA. In addition, 150 ng of biotinylated BMP2 cDNA was added to remove BMP-2-transcripts from the subtracted library. The mixture was denatured and hybridized for 20 h at 68 °C. After hybridization biotinylated molecules were removed from the mixture with Dynabeads M-280 Streptavidin (DYNAL). The resulting subtracted cDNA was ethanol precipitated and hybridized with another 15 μ g of biotinylated control cDNA and the cycle was repeated. The resulting cDNA was amplified by PCR (15 cycles) using the target-library specific 21-mer oligonucleotide as primer. In total six hybridization/subtraction cycles were performed. The resulting cDNAs were cleaved with *Hind* III, ligated into the eukaryotic expression vector pMBC-2T-fl and transformed into competent *E. coli* SURE cells. The colonies were transferred to nitrocellulose and a plus-minus-screening was performed using the subtractive library and the control/driver library as probes. ~30% of the clones proved to be differentially expressed and were further characterized by DNA sequencing and northern analyses.

Mice and RNA-*in situ*-hybridization

Embryos were isolated from pregnant NMRI mice at the developmental stages indicated in the text. The day of plug detection was considered to be day 0.5 post conceptionem (dpc). The embryos were fixed overnight with 4% paraformaldehyde in PBS at 4°C. For radioactive RNA-*in situ*-hybridization fixed mouse embryos and tissues were prepared for cryostat sectioning as described previously (Bächner et al., 1993). 10 μ m cryosections were mounted on 3-aminopropyltriethoxysilane coated slides. Antisense and sense riboprobes were generated by RNA *in vitro* transcription with [³⁵S]-dUTP to a specific activity of >10⁹ dpm/ μ g. Probe length was reduced to 150 - 200 nucleotides by alkaline hydrolysis (Cox et al., 1984). The slides were prehybridized at 54°C in a solution containing 50% formamide, 10% dextrane sulfate, 0.3 M NaCl, 10 mM Tris, 10 mM sodium phosphate pH 6.8, 20 mM dithiothreitol, 0.2x Denhardt's reagent, 0.1% Triton X-100, 1.25 mg/ml yeast RNA, and "cold" 0.1 mM [S]-dUTP. For hybridization, 80000 dpm/ μ l [³⁵S]-dUTP labelled probe RNA was added to the hybridization mix, and the hybridization was continued at 54°C for 16 h in a humid chamber. The slides were washed in hybridization salt solution to which dithiothreitol was added. After RNase A digestion (40 μ g/ml) the slides were washed for 30 min at 37°C with 2x SSC, 0.1% (w/v) SDS and 30 min with 0.1x SSC and dehydrated by increasing concentrations of ethanol. The slides were coated with Ilford K5 photoemulsion for autoradiography. After 1 to 4 weeks of exposure at 4°C, depending on cDNA analysed, the slides were developed and stained with Giemsa solution. The embryos and sections were analyzed with bright- and dark-field illumination with a Zeiss SV11 stereo-microscope and an Zeiss Axiophot microscope and photographed using Kodak Ektachrome 320T or Agfa Ortho 25 film. Figures were prepared using a Polaroid SlideScanner together with Adobe Photoshop and Adobe PageMaker software with a Windows NT-computer device.

Results

Expression of BMPs in C3H10T $\frac{1}{2}$ mesenchymal precursor cells

To compare the *in vitro* characteristics of BMP-mediated differentiation with other members of the BMP-family the cDNAs encoding the human version for BMP2-7 (without BMP3) were integrated into an expression vector (Ahrens et al., 1993) and transfected into C3H10T $\frac{1}{2}$ mesenchymal precursor cells. 5.000-10.000 transfectants per individual member of the BMP-family were pooled and investigated further. After reaching confluence all BMP-transfected cells started to grow in multilayer and extensive matrix production accompanies a development into three distinct mesenchymal lineages: the adipogenic, the osteogenic and the chondrogenic (Fig. 1) (Ahrens et al., 1993; Wang et al., 1993).

Although in general the differentiation in osteoblasts, chondrocytes and adipocytes was observed for all BMPs investigated on the basis of morphological and histological criteria (Fig. 1), the ability of the various BMP-expressing C3H10T $\frac{1}{2}$ cells to undergo mesenchymal development within a two week cultivation period was different among the various members of the BMP-family (Table I). In all cases, the level of BMP-specific mRNA was comparable for the various BMP-expressing C3H10T $\frac{1}{2}$ cells. The mRNA encoding the BMPs are, in general, expressed at the highest level in actively proliferating cells up to few days after reaching confluence (Fig. 2). Secreted BMPs are difficult to monitor in the supernatant of C3H10T $\frac{1}{2}$ cells. We could confirm in another study by western analyses that BMPs such as BMP2 and BMP4 can only marginally be monitored in the supernatant, indicating that they are efficiently bound by extracellular matrix (Wang et al., 1993).

Isolation of BMP-2 induced cDNAs from the mesenchymal progenitor cell line C3H10T $\frac{1}{2}$

We employed the subtractive cloning procedure outlined in Materials and Methods to C3H10T $\frac{1}{2}$ cells expressing recombinant BMP2 at 8 day post-confluence. At this stage the first morphologically distinct phenotypes of the various mesenchymal lineages are monitored and, therefore, we should be able to isolate cDNAs which are involved in the

determination of these lineages as well as cDNAs which could play a role in the morphological manifestation of the mesenchymal cell types. The subtractive cloning strategy involves two different PCR-primer sets for the PCR-amplification of the driver and the target library and the addition of biotinylated BMP2 cDNA to the target library to prevent the selection of cDNAs originating from recombinant BMP2-transcripts. After the subtraction procedure, *E. coli* colonies harboring the cloned cDNA fragments were transferred to nitrocellulose, and a plus-minus-screening was performed using the amplified cDNAs from the subtractive library and the control/driver library as probes. Roughly 10.000 colonies were screened and about 30 % of the clones proved to be differentially expressed. Of these 200 were further characterized by northern analyses and DNA sequencing.

Partially or complete sequencing of 21 BMP-regulated cDNAs revealed that 20 were related to known sequences and one was not (29 A) (Table II). About 50 % of the cDNAs consisted of four different sequence types (Table II). The others were represented between 1 - 4% in the subtracted library. Among the known sequences were collagens, several enzymes of the glycolytic pathway, cystatin C, vimentin, basigin, tropoelastin, migration inhibitory factor (MIF), osteopontin, lipoprotein lipase, the heatshock protein HSP-47, autotaxin (ATX) as well as one member of the family of the CCAAT-binding family of transcription factors, C/EBP α . Of the former some were not previously recognized as BMP-upregulated genes, namely C/EBP α , HSP-47, MIF, cystatin C, basigin, vimentin, G0S2, tropoelastin, autotaxin (ATX) and 29A. The expression profile of these genes was characterized by northern analyses in parental and recombinant BMP2 expressing C3H10T $\frac{1}{2}$ cells (Fig. 2) which shows that the genes isolated by subtractive cloning were expressed predominantly at middle to late cultivation stages (e. g. autotaxin, G0S2, tropoelastin) but some BMP-regulated mRNAs are regulated already early on in recombinant BMP-expressing cells like e. g. HSP47, vimentin and especially the undescribed cDNA 29A which encodes a putative secreted factor (see below) (Fig. 2).

BMP-regulated gene 29A isolated from C3H10T $\frac{1}{2}$ cells is involved in the development of the osteo- /chondrogenic lineage.

29A has been isolated as a 250 bp cDNA fragment hybridizing to a BMP-upregulated 1.8 kb mRNA in C3H10T $\frac{1}{2}$ cells (Fig. 2). The cDNA sequence has been completed by the RACE technology and confirmed by cDNA cloning. It contains several open reading frames. Already the first translational start site obeys the Kozak rules and is followed by an open reading frame coding for a protein with a M_w 22 kDa (Fig. 6a). 29A seems as a secreted factor indicated by the N-terminal hydrophobic stretch of 30 amino acid residues. The putative signal-sequence cleavage-site has been located between amino acid positions 23-24 (Signal-program at ExPASy-Tools: ISREC, Lausanne). The protein sequence does not exhibit a significant homology to known sequences. During murine development 29A expression is detected in a variety of mesodermal tissues. Enhanced expression is first detected in presumptive bone-forming centers of vertebrae at 12.5 dpc (Fig. 7e, f) and later gets restricted to the perichondrium of the forming vertebrae (Fig. 7c, d, g, h). In midgestation development expression is further detected in the ribs, tooth bud and forming vibrissae (Fig. 3f; 5d, 7c, d). During limb development enhanced expression gets restricted to the perichondrium and connective tissue sheet of the forming metatarsals and phalanges (Fig. 4d). In late tooth development 29A expression is restricted to the outer enamel epithelium and the ameloblasts (Fig. 5d). 29A is also expressed in primary murine osteoblasts from 5-day old mice suggesting a potential regulatory role for this protein in bone growth (Fig. 6c).

Discussion

BMP2 upregulated cDNAs in C3H10T $\frac{1}{2}$ cells

A new finding was the unknown BMP2-upregulated gene 29A which is expressed early in C3H10T $\frac{1}{2}$ cells and primary osteoblasts. In the latter cells its expression increases at late stages of the development *in vitro* which is comparable with many osteogenic marker genes (Fig. 6c). The putative secreted factor 29A could be one of the many growth factors which influence the initiation and the manifestation of the bone and/or cartilage phenotype. Such factors exist in a wide variety involving very different factors like e. g. insulin-like growth factors (IGFs), parathyroid hormone related protein (PTHrP) and Indian hedgehog (IHH). At present the role of 29A is determined in several differentiation systems. Such studies also show that a homologous factor is found in the human system.

References

- Ahrens, M., T. Ankenbauer, D. Schroder, A. Hollnagel, H. Mayer, and G. Gross. 1993. Expression of Human Bone Morphogenetic Proteins - 2 or - 4 in Murine Mesenchymal Progenitor C3H10T $\frac{1}{2}$ Cells Induces Differentiation into Distinct Mesenchymal Cell Lineages. *DNA Cell Biol.* 12:871-880.
- Bächner, D., A. Manca, P. Steinbach, D. Wöhrle, W. Just, W. Vogel, H. Hameister, and A. Poustka. 1993. Enhanced expression of the murine FMR1 gene during germ cell proliferation suggests a special function in both the male and

female gonad. *Hum. Mol. Genet.* 2:2043-2050.

Blessing, M., L. B. Nanney, L. E. King, C. M. Jones, and B. L. M. Hogan. 1993. Transgenic mice as a model to study the role of TGF- β -related molecules in hair follicles. *Genes Dev.* 7:204-215.

Cox, K. H., D. V. DeLeon, L. M. Angerer, and R. C. Angerer. 1984. Detection of mRNAs in sea urchin embryos by in situ hybridization using asymmetric RNA probes. *Dev. Biol.* 101:485-502.

Ferguson, E. L. and K. V. Anderson. 1992. Localized enhancement and repression of the activity of the TGF- β family member, *decapentaplegic*, is necessary for dorsal-ventral pattern formation in the *Drosophila* embryo. *Development* 114:583-597.

Frohmann, M. A. 1990. Rapid amplification of cDNA ends (RACE): user-friendly cDNA cloning. *Amplifications* 5:11-15.

Hofmann, C., G. B. Luo, R. Balling, and G. Karsenty. 1996. Analysis of limb patterning in BMP-7-deficient mice. *Dev. Genet.* 19:43-50.

Irish, V. F. and W. M. Gelbart. 1987. The decapentaplegic gene is required for dorsal-ventral patterning of the *Drosophila* embryo. *Genes Dev.* 1:868-879.

Karsenty, G., G. B. Luo, C. Hofmann, and A. Bradley. 1996. BMP 7 is required for nephrogenesis, eye development, and skeletal patterning. *Ann. NY Acad. Sci.* 785:98-107.

Lyons, K. M., J. L. Graycar, A. Lee, S. Hashmi, P. B. Lindquist, E. Y. Chen, B. L. M. Hogan, and R. Derynck. 1989a. Vgr-1, a mammalian gene related to xenopus Bg-1 is a member of the transforming growth factor β gene superfamily. *Proc. Natl. Acad. Sci. USA* 86:4554-4558.

Lyons, K. M., R. W. Pelton, and B. L. M. Hogan. 1989b. Patterns of expression of murine Vgr-1 and BMP-2a RNA suggest that transforming growth factor- β -like genes coordinately regulate aspects of embryonic development. *Genes Dev.* 3:1657-1668.

Mishina, Y., A. Suzuki, N. Ueno, and R. R. Behringer. 1995. Bmpr encodes a type I bone morphogenetic protein receptor that is essential for gastrulation during mouse embryogenesis. *Genes Dev.* 9:3027-3037.

Owen, T. A., M. Aronow, V. Shalhoub, L. M. Barone, L. Wilming, M. S. Tassinari, M. B. Kennedy, S. Pockwinse, J. B. Lian, and G. S. Stein. 1990. Progressive Development of the rat osteoblast phenotype in vitro, reciprocal relationships in the expression of genes associated with osteoblast proliferation and differentiation during formation of the bone extracellular matrix. *J. Cell. Physiol.* 143:420-430.

Reznikoff, C. A., D. W. Brankow, and C. Heidelberger. 1973. Establishment and characterization of a cloned cell line of C3H mouse embryo cells sensitive to postconfluence inhibition of division. *Cancer Res.* 33:3231-3238.

Taylor, S. M. and P. A. Jones. 1979. Multiple new phenotypes induced in 10T $\frac{1}{2}$ and 3T3 cells treated with 5-azacytidine. *Cell* 17:771-779.

Wang, E. A., D. I. Israel, S. Kelly, and D. P. Luxenberg. 1993. Bone-Morphogenetic protein causes commitment and differentiation in C3H10T $\frac{1}{2}$ and 3T3 cells. *Growth Factors* 9:57-71.

Weeks, D. L. and D. A. Melton. 1987. A maternal mRNA localized to the vegetal hemisphere in xenopus eggs codes for a growth factor related to TGF β . *Cell* 51:861-867.

Winner, G., M. Blessing, P. A. Labosky, and B. L. M. Hogan. 1995. Bone-morphogenetic protein-4 is required for mesoderm formation and patterning in the mouse. *Genes Dev.* 9:2105-2116.

Wozney, J. M. 1989 Bone morphogenetic proteins. *Prog. Growth Factor Res.* 1:267-280.

Legend to Tables

Table I. Differentiation potential of C3H10T $\frac{1}{2}$ cells stably expressing members of the BMP-family. Numbers represent cells/cm². Cells were grown in 4.5 cm² wells. The total number of cells was determined in a coulter counter after treatment of the dense adherent cellular layer with collagenase. The values represent mean values of three independent cultivations. The number of osteoblast-like cells were evaluated by assessing the number of alkaline phosphatase positive colonies. The number of chondroblastic cells was determined after staining with Alcian Blue. Adipocytes were identified morphologically or stained with Oil Red O. The multilayer growth obstructed exact evaluation of the mesenchymal lineages at later stages of cultivation. Therefore, the number of osteoblast-like and chondroblastic cells represent approximated values of three independent cultivations. The number of adipocytes which heavily covered the cells was not determined.

Table II. BMP-regulated genes isolated by subtractive cloning from recombinant BMP2-expressing C3H10T $\frac{1}{2}$ cells. Subtractive cloning is detailed in Materials and Methods. ~200 BMP-upregulated cDNAs after subtractive cloning were investigated in northern analyses and by sequencing. The percentages indicate the representation of the respective cDNAs in the subtractive cDNA library. The other cDNAs are represented between ~1%-4% in the library. * indicate members of the C/EBP-family which have not been detected by subtractive library but were found to be upregulated by BMP during the course of this study (see also Fig.4)

Legends to Figures:

Fig. 1. Histological analysis of mesenchymal development in C3H10T $\frac{1}{2}$ cells. a. C3H10T $\frac{1}{2}$ cells stably transfected with the expression vector pMT7T3 (10 days after reaching confluence) b. Alkaline phosphatase positive osteoblast-like cells in C3H10T $\frac{1}{2}$ cells stably expressing BMP2 (10 days post-confluence) c. Alcian blue positive, chondrocyte-like cells in C3H10T $\frac{1}{2}$ cells stably expressing BMP7 (12 days post-confluence). d. oil-droplet filled adipocyte-like cells in C3H10T $\frac{1}{2}$ cells stably expressing BMP4 (12 days post-confluence).

Fig. 2 Northern analyses of BMP2-regulated genes isolated by subtractive cloning from C3H10T $\frac{1}{2}$ cells. Northern analyses were performed as described in Fig. 2 and Materials and Methods.

Fig. 3. Expression of BMP-regulated genes in murine midgestation development. Bright (a) and dark field (b-f) images of consecutive parasagittal cryosections of a 16.5 dpc mouse embryo hybridized with a HSP47 (b), collagen α_1 (I), GOS2 (d), ATX (e), and 29A (f) antisense riboprobe are shown. HSP47 and collagen (I) are coexpressed in many cartilage tissues, for example the ribs and cranial bones (b,c). Expression of GOS2 is in contrast restricted to forming brown adipose tissue (arrow in d), a lower level of expression is also visible in the liver (d). ATX expression is most prominent in the mesenchyme surrounding the forming vibrissae, the choroid plexus epithelia of the IVth. (arrow in e) and Ist (arrowhead in e) ventricle of the brain. Enhanced expression of ATX is in addition visible in different cartilage tissues, for examples within the hind limb, ribs, and cranial bones (e). Expression of 29A is most prominent in the hind limb, ribs (arrow in f), and epithelia of the brain (arrowhead in f). cp, choroid plexus; co, cochlea; li, liver; lu, lung; me, mesencephalon; rb, rib; te, telencephalon; vi, vibrissae. Bar, 1 mm.

Fig. 4. Expression of BMP-regulated genes within 16.5 dpc hind limb. Higher magnifications of the hind limb of Fig. 5 are shown. Whereas HSP47 expression within the developing limb is most prominent in the cartilage of the forming bones (b), expression of ATX is restricted to the joint regions of the forming bones (c), and expression of 29A most prominent in the perichondrium and adjacent connective tissue sheet of the metatarsals and phalanges (d). cal, calcaneum; cun, cuneiforme; mta, metatarsale; pha, phalange; sk, skin; tal, talus; tib, tibia. Bar, 100 μ m.

Fig. 5. Expression of BMP-regulated genes within 18.5 dpc lower tooth bud. Consecutive cryosections through the lower tooth bud region of a 18.5 dpc mouse embryo hybridized with a HSP47 (b), ATX (c), and 29A (d) antisense riboprobe are shown. Expression of HSP47 is most prominent in odontoblasts and the alveolar bone (b). In contrast ATX expression is enhanced in the stratum intermedium layer, separating the ameloblast layer from the stellate reticulum (c). A lower level of expression is visible in mesenchymal tissue adjacent to tooth bud (c). Expression of 29A is most prominent in the outer enamel layer, but also visible in the ameloblasts (d). alb, alveolar bone; amb, ameloblasts; dep, dental papillae; mes, mesenchyme; odb, odontoblasts; oen, outer enamel epithelium; str, stratum intermedium. Bar, 100 μ m.

Fig. 6. Structure of the 29A cDNA and its expression profile in primary osteoblasts. a. Sequence of the 29A cDNA. The first open region reading frame gives rise to a putative secreted 22 kDa protein. b. Features of the BMP-upregulated cDNAs ATX and 29A are schematically represented. c. 29A is expressed *in vitro* during the osteoblast developmental sequence in cultivated primary osteoblasts isolated from 5 day old mice

Fig. 7. Expression of 29A in mouse development. Parasagittal cryosections of 12.5 (e,f), 14.5 (a,b,g,h) and 16.5 (c,d,i,k) mouse embryos hybridized with a 29A antisense riboprobe are shown. Enhanced expression of 29A is visible in preanilage condensations of the vertebrae at 12.5 dpc (e,f). At 14.5 dpc, expression is restricted to the perichondrium of the forming vertebrae (g,h) but also visible within cartilage of the hind limb buds and ribs (b), and the faciae surrounding the spinal ganglia (arrow in b and h). Enhanced expression is in addition visible at the forming vibrissae (a,b). At 16.5 dpc expression remains enhanced in the perichondrium of the vertebrae within the tail (c,d), and the perichondrium and connective tissue sheet of the metatarsals and phalanges of the hind limb (i,k), but also at the vibrissae (c,d). bl, bladder; cts, connective tissue sheet; gt, gut; hlb, hind limb bud; li, liver; mta, metatarsale; my, myelencephalon; per, perichondrium; pha, phalanges; sga, spinal ganglia; sk, skin; ta, tail; te, telencephalon; ve, vertebra; vi, vibrissae. Bar. ad. 1 mm; e-k, 100 μ m.

Table I

Days after confluence		0	4	7	9	11
BMP-2	Osteoblastic	0	0-1	20	220	1100
	Chondroblastic	0	0	20	45	45
	Adipocytic	0	0	0	covered	covered
	Total number of cells	0.19×10^6	0.53×10^6	0.66×10^6	0.66×10^6	0.68×10^6
BMP-4	Osteoblastic	0	0	2	20	120
	Chondroblastic	0	0	0	2	2
	Adipocytic	0	0	0	covered	covered
	Total number of cells	0.16×10^6	0.26×10^6	0.42×10^6	0.44×10^6	0.48×10^6
BMP-5	Osteoblastic	0	0	1	5	20
	Chondroblastic	0	0	5	20	100
	Adipocytic	0	0	0	100	500
	Total number of cells	0.13×10^6	0.22×10^6	0.23×10^6	0.22×10^6	0.22×10^6
BMP-6	Osteoblastic	0	0-2	110	1100	2250
	Chondroblastic	0	0	0	120	210
	Adipocytic	0	0	0	covered	covered
	Total number of cells	0.14×10^6	0.4×10^6	0.37×10^6	0.42×10^6	0.48×10^6
BMP-7	Osteoblastic	0	0	2	10	20
	Chondroblastic	0	0	50	100	100
	Adipocytic	0	0	0	covered	covered
	Total number of cells	0.21×10^6	0.17×10^6	0.37×10^6	0.35×10^6	0.36×10^6
10T $\frac{1}{2}$	Total number of cells	0.15×10^6	0.15×10^6	0.17×10^6	0.17×10^6	0.18×10^6

Table II

cDNAs isolated with homology to:	Representation in the subtracted library
ApoE Autotaxin (ATX) Basigin C/EBP α C/EBP β * C/EBP δ * Collagen (I) Collagen (II) Collagen (III) Cystatin C	8%

Table II (continued)

cDNAs isolated with homology to:	Representation in the subtracted library
Enolase (2-phospho-D-Glycerate-Hydrolase)	20%
G0S2	
Glycerolaldehydedehydrogenase (GAPDH) / Uracyl-DNA-Glykosylase (UDG)	10%
HSP47	
Lipoproteinlipase	
Migration Inhibitory Factor (MIF)	10%
Osteopontin	
Phosphofructokinase (PFK)	
Phosphoglyceratekinase (PGAM)	
Pyruvatekinase (PK)	
Tropoelastin	
Vimentin	
29A	

Claims

1. ssDNA

(a) having a sequence of from position 1 to 1753 according to Fig. 6a or

(b) having a truncated sequence compared with the ssDNA according to (a), wherein its nucleotides are within the range of from position 1 to 1753 according to Fig. 6a or

(c1) having a sequence which differs from that of the ssDNA according to (a) but has the same number of nucleotides or

(c2) having a sequence which differs from that of the ssDNA according to (b) but has the same number of nucleotides

wherein the ssDNA according to (c1) or (c2) is hybridizable with that according to (a) and (b), respectively.

2. A ssDNA according to claim 1

(a) having a sequence of from position 417 to 1022 or

(b) having a sequence of from position 489 to 1022.

3. ssDNA according to claim 1 (c1) or claim 1 (c2) or having a sequence which differs from that of a ssDNA according to claim 2 but has the same number of nucleotides, each being hybridizable at a temperature of at least 25 °C and a 1M sodium chloride concentration.

4. A ssDNA, characterized in that it is complementary to a ssDNA according to any of the preceding claims.

5. dsDNA, consisting of a ssDNA according to any of the preceding claims and its complementary strand.

6. Vector, comprising a dsDNA according to claim 5.

7. An expression product of a vector according to claim 6.

8. A cell comprising a vector according to claim 6.

9. Use of a vector according to claim 6 or an expression product according to claim 7 for a therapeutical regulation of bone growth, cartilage growth, tissue regeneration, tissue remodelling or wound healing.

10. Use of a vector according to claim 6 or an expression product according to claim 7 or a cell according to claim 8 for the detection of agonists and/or antagonists of bone, cartilage or tissue growth regulation.

11. Use of a ssDNA according to any of claims 1 to 4, a dsDNA according to claim 5 or an expression product according to claim 7 for diagnostic tests.




Fig. 1






Fig. 2



Fy. 3



Fj.4



HSP47



29A



16.5 dpc, hind limb



ATX

Fig. 5


b

29A

Sag uipryside




Autotaxin (ATX)



C

0 4 8 14 20 [days]



Ty. 6

a

(a)
enlarged

CGTCCCTGGCCTTGCCCTAACTCTTCTGTGGTCTGTAACATTACCTGTGAATTTCCCA 60
 GCCGAAACGGTGTGGGGCAAGAACTTCTTGTAAAACCTCCACCCCTTGGACTCTCC 120
 ACAGCCCTCTCACCCTCCCAATCTTCTGAGACGCTTTTACCTCTCCGCCAGAGCAGAG 180
 TTTATCTTTTTTTCTTTTGCTTTTTTTCTTTTCTCCCATTTTCTCGCCCTGICC 240
 TTTACATCTGAAAGGAGATCAGTTCAAGAGTGACCAGGTGGACGCTCCTTTTCTTAT 300
 TTAGTTTATTATTGTTTGGGGAGTTTCTTTCTATCTTTTTTAATTCCTGTCCGGGGA 360
 GTTTTGTCCACCGCTCCTACCACCTCCCCCTGTACCCCGCTCCGCCGCGGAGGATGG 420
 ValTrpLysTrpLeuGlyAlaLeuValValPheProLeuGlnMetIleTyrLeuValThr
 TGTGGAAATGGCTGGGCGCGCTGGTAGTGTCCCTCTGCAATGATCTATTTGGTAACCA 480
 LysAlaAlaValGlyMetValLeuProProLysLeuArgAspLeuSerArgGluSerVal
 AAGCAGCGGTGGGAATGGTGTGCCCCCAAGCTTCGGGACTTGTGCGGGAGTCAGTCC 540
 LeuIleThrGlyGlyGlyArgGlyIleGlyArgHisLeuAlaArgGluPheAlaGluArg
 TCATCACCAGCGGTGGGAGAGGCATCGGACGCCACCTCGCTCGGGAGTTCGCAGAGCGTG 600
 GlyAlaArgLysIleValLeuTrpGlyArgThrGluLysCysLeuLysGluThrThrGlu
 GCGCCAGAAAGATTGTTCTCTGGGGCGGACTGAAAAATGCCCAAGGAGACGACAGAGG 660
 GluIleArgGlnMetGlyThrGluCysHisTyrPhelleCysAspValGlyAsnArgGlu
 AGATTCGGCAGATGGGCACAGAGTGCCACTACTTCATCTGTGACGTGGGCAACCGGAAG 720
 GluValTyrGlnMetAlaLysAlaValArgGluLysValGlyAspIleThrIleLeuVal
 AGGTGTACCAGATGGCCAAAGCTGTCCGAGAGAAGGTGGGTGACATCACCATCCTGGTGA 780
 AsnAsnAlaAlaValValHisGlyLysSerLeuMetAspSerAspAspAlaLeuLeu
 ACAATGCCGCTGTGGTCCATGGAAGGCTTGATGGACAGTGACGATGATGCCCTCCTCA 840
 LysSerGlnHisValAsnThrLeuGlyGlnPheTrpThrThrLysAlaPheLeuProArg
 AGTCCAGCATGTCAACACCTGGGCCAATTCTGGACCACCAAGGCCTTTTGCCACGTA 900
 MetLeuGluLeuGlnAsnGlyHisIleValCysLeuAsnSerValLeuCysHisValSer
 TGCTGGAATCCAGAACGGCCATATTGTGTGCTCAATTCGTGCTTTGCCATGTCAGCC 960
 HisProTrpGlnSerIleAspTyrCysThrSerLysAlaSerAlaSerProSerTrpArg
 ATCCCTGGCAGTCCATCGACTACTGCACGTCAAAGCGTCAGCTTCGCTTCATGGAGAG 1020
 Ala
 CCTGACCTTGGGGCTGTGGACTGCTGCTGGTGTGACGCCACCACCGTTCTGCCCTTTCA 1080
 CACCAGCACCGAGATGTTCCAGGGCATGAGAGTCAGGTTTCCCAACCTCTTCCGCCACT 1140
 GAGGCCAGGGACAGTAGCCCGGAGACGGTAGATGCTGTGCAACAAACCAGGCCCTTCTC 1200
 TTICACCCGTTGGACCATGAATATTCTCATTGTCTTGAAAGCATACTCCACAAAGCTGCA 1260
 CTGGAGGAGATTACAGGTTCTCGGGGACTTACACCTGTATGAACACCTTTAAGGGGAGG 1320
 ACATAGAGGACAGGAGGAAGACACCTGAGGAGCTATGGAGCCTGAGGGGAGCCACAGC 1380
 AGCCGGGCACACAATCCTGTGCCTGTGCATTAGCACATCTGCTGGGTGAACAGGACTGTT 1440
 CTTGTCCCCAGGGAGATTTTTTCAGCTCCCCAGGTCACTCCAGGACCTTTGTGCAAGAC 1500
 TGATGGGTTTAACTCTGACCCCATGGAGGCAAGAAGCCGGCAGCCCCCAACAACCTTG 1560
 TACATTTCTCATTTCTGAGCGTTTGTGATGAAATCGCTTCTCCAGTCTAACCCGCCTAAT 1620
 GTGCATCTACTATTTCAGGAGAGTCTGCTCCAGACACTCTGCCTTTCCCTCCAAAACC 1680
 CTCTCACTCCAGCTCGTGCAAACTGGTTACACAGCAGAAACGAAAAATAAGAGGTGCG 1740
 TTTTCAAAAAAAA





Fig. 7



(19)

Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 0 890 639 A3

(12)

EUROPEAN PATENT APPLICATION

(88) Date of publication A3:
10.10.2001 Bulletin 2001/41

(51) Int Cl.7: **C12N 15/12, C07K 14/47,
C07K 14/51, A61K 38/17**

(43) Date of publication A2:
13.01.1999 Bulletin 1999/02

(21) Application number: **98112742.6**

(22) Date of filing: **09.07.1998**

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE**
Designated Extension States:
AL LT LV MK RO SI

(30) Priority: **09.07.1997 EP 97111602**

(71) Applicant: **Gesellschaft für Biotechnologische
Forschung mbH (GBF)**
38124 Braunschweig (DE)

(72) Inventors:
• **Ahrens, Marion**
38124 Braunschweig (DE)
• **Bächner, Dietmar**
38124 Braunschweig (DE)

- **Hoffmann, Andrea**
38124 Braunschweig (DE)
- **Lauber, Jörg**
38124 Braunschweig (DE)
- **Steinert, Peter**
38124 Braunschweig (DE)
- **Flohé, Leopold**
38124 Braunschweig (DE)
- **Gross, Claus-Gerhard**
38124 Braunschweig (DE)

(74) Representative: **Boeters, Hans Dietrich, Dr. et al**
Patentanwälte Boeters & Bauer,
Bereiteranger 15
81541 München (DE)

(54) **BMP2-induced cDNA and its use**

(57) The recombinant expression of human BMP2, BMP4 - BMP7 in murine mesenchymal C3H10T½ progenitors mediates differentiation into three mesenchymal lineages in different efficiencies: the osteogenic, the chondrogenic and the adipogenic lineage. This developmental *in vitro* model was used to identify and to char-

acterize cDNAs involved during the manifestation of these lineages *in vivo*. By subtractive cloning an as yet undescribed cDNA, **29A** has been cloned which encodes a putative secreted factor which is expressed in developing osteo- /chondrogenic tissues of vertebrae, ribs, tooth and the limb bud.

EP 0 890 639 A3



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 98 11 2742

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
X	<p>DATABASE NUCLEOTIDE AND PROTEIN 'Online! EBI, UK1 March 1996 (1996-03-01) "Mus musculus mRNA for retinal short-chain dehydrogenase/reductase retinal short-chain dehydrogenase/reductase; retSDR1 gene" Database accession no. X95281 XP002173431 * abstract *</p>	1-4	<p>C12N15/12 C07K14/47 C07K14/51 A61K38/17</p>
X	<p>DATABASE NUCLEOTIDE AND PROTEIN 'Online! EBI, UK23 June 1996 (1996-06-23) "Soares mouse embryo NbME13.5 14.5 Mus musculus cDNA clone IMAGE:390287 5' similar to SW:YOXD_BACSU p14802 HYPOTHETICAL 25.3 KD PROTEIN IN RTP 5'REGION, mRNA sequence, EST" Database accession no. W74923 XP002173432 * abstract *</p>	1-4	
X	<p>DATABASE NUCLEOTIDE AND PROTEIN 'Online! EBI, UK2 April 1997 (1997-04-02) "Barstead MPLRB1 Mus musculus cDNA clone IMAGE:766482 5' similar to WP:T11F9.11 CE06423 3-OXOACYL-'ACYL-CARRIER PROTEIN! REDUCTASE, mRNA sequence, EST" Database accession no. AA275737 XP002173433 * abstract *</p>	1-4	<p>TECHNICAL FIELDS SEARCHED (Int.Cl.6) C12N A61K</p>
The present search report has been drawn up for all claims			
Place of search BERLIN		Date of completion of the search 31 July 2001	Examiner Mateo Rosell, A.M.
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background C: non-written disclosure F: intermediate document</p>		<p>T: theory or principle underlying the invention E: earlier patent document, but published on, or after the filing date D: document cited in the application I: document cited for other reasons S: member of the same patent family, corresponding document</p>	



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 98 11 2742

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
D.Y	AHRENS M ET AL: "Expression of human bone morphogenetic proteins-2 or -4 in murine mesenchymal progenitor C3H10T1/2 cells induces differentiation into distinct mesenchymal cell lineages" DNA AND CELL BIOLOGY, US, NEW YORK, NY, vol. 12, no. 10, 1 December 1993 (1993-12-01), pages 871-880, XP002087786 ISSN: 1044-5498 * the whole document *	1-11	TECHNICAL FIELDS SEARCHED (Int.Cl.6)
Y	WO 93 05149 A (HUTCHINSON FRED CANCER RES) 18 March 1993 (1993-03-18) * page 4, line 26 - page 6, line 16 * * page 21, line 19 - page 23, line 7 * * page 30, line 36 - page 36, line 10 *	1-11	
A	WO 96 38590 A (OSTEOSCREEN INC ; HARRIS STEPHEN E (US); MUNDY GREGORY R (US); GHOS) 5 December 1996 (1996-12-05) * page 2, line 22 - page 3, line 11; examples 1-5 *	1-11	
A	HARA E ET AL: "DNA-DNA SUBTRACTIVE CDNA CLONING USING OLIGO(DT)30-LATEX AND PCR: IDENTIFICATION OF CELLULAR GENES WHICH ARE OVEREXPRESSED IN SENESCENT HUMAN DIPLOID FIBROBLASTS" ANALYTICAL BIOCHEMISTRY, US, ACADEMIC PRESS, SAN DIEGO, CA, vol. 214, no. 1, 1 October 1993 (1993-10-01), pages 58-64, XP000396227 ISSN: 0003-2697 * the whole document *	1	
The present search report has been drawn up for all claims			
Place of search:		Date of completion of the search	Examiner
BERLIN		31 July 2001	Mateo Rosell, A.M.
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background O: non-written disclosure P: intermediate document</p> <p>1: theory or principle underlying the invention E: earlier patent document, but published on, or after the filing date D: document cited in the application L: document cited for other reasons 8: member of the same patent family corresponding document</p>			



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 98 11 2742

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document, with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
T	<p>BAECHNER DIETMAR ET AL: "Bmp-2 downstream targets in mesenchymal development identified by subtractive cloning from recombinant mesenchymal progenitors (C3H10T1/2)."</p> <p>DEVELOPMENTAL DYNAMICS, vol. 213, no. 4, December 1998 (1998-12), pages 398-411. XP001015326</p> <p>ISSN: 1058-8388</p> <p>* the whole document *</p> <p>-----</p>	1-11	
			TECHNICAL FIELDS SEARCHED (Int.Cl.6)
The present search report has been drawn up for all claims			
Place of search		Date of completion of the search	Examiner
BERLIN		31 July 2001	Mateo Rosell, A.M.
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X: particularly relevant if taken alone</p> <p>Y: particularly relevant if combined with another document of the same category</p> <p>A: technological background</p> <p>C: non-written disclosure</p> <p>P: intermediate document</p> <p>1: theory or principle underlying the invention</p> <p>2: earlier patent document, but published on, or after the filing date</p> <p>3: document cited in the application</p> <p>4: document cited for other reasons</p> <p>5:</p> <p>6: member of the same patent family, corresponding document</p>			

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 98 11 2742

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

31-07-2001

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
WO 9305149	A	18-03-1993	AU	2571892 A	05-04-1993
WO 9638590	A	05-12-1996	US	6083690 A	04-07-2000
			AU	6377996 A	18-12-1996
			CA	2221621 A	05-12-1996
			EP	0832297 A	01-04-1998
			JP	11506009 T	02-06-1999

EP 0 890 639 A3

For more details about this annex, see Official Journal of the European Patent Office, No. 12/82